## Contents

1. Abstract	1
2. Introduction	2
2.1 Formation of a polarized Golgi	4
2.2 Intra-Golgi transport	7
2.3 The Golgi as a platform to coordinate signalling cascades	9
2.4 Lipid-enriched microdomains	10
2.5 Golgi-derived lipid rafts	12
2.6 GPI-anchored proteins	16
2.7 Trafficking of GPI-anchored proteins	19
2.8 Purpose of the thesis	23
3. Results	24
3.1 Isolation of GREG	24
3.1.1 Cloning of the cDNA	25
3.1.2 Characteristics of GREG	29
3.1.3 GREG has a luminal orientation	29
3.1.4 GREG is a glycoprotein	30
3.1.5 Confirmation of the predicted ORF	32
3.2 The coiled-coil domain is involved in protein-protein interactions	34
3.3 GREG is homologous to human BST-2	34
3.4 GREG is a GPI-anchored protein	36
3.4.1 Aerolysin detects a protein in GICs similar to GREG	37
3.4.2 Expression of FLAG-tagged GREG in mammalian cells	38
3.4.3 GPI linkage of GREG	40
3.5 GREG resides at the Golgi complex	42
3.5.1 Immunolocalization of the tagged GREG	42
3.5.2 Biochemical localization of the native and tagged GREG	44
3.6 Mapping the Golgi targeting domain of GREG	46
3.6.1 The C-terminal domain of GREG contains	
a Golgi targeting signal	47
3.6.2 EQ repeat is not sufficient for Golgi targeting	
of a soluble protein	49
3.6.3 Is the EQ repeat a universal Golgi targeting signal?	49

	3.7 GREG does not recycle between phagosomes and the Golgi complex	51
	3.8 GREG is essential in maintenance of the Golgi integrity	52
	3.8.1 Blockade of GREG expression vesiculates the Golgi	53
	3.8.2 A dominant negative mutant of GREG	54
	3.8.3 Membrane attachment is not enough for GREG to function	55
	3.9 Recruitment into GICs is required for GREG to function	57
	3.10 Golgi fragmentation by mutant GREG is independent	
	of apoptosis and microtubule integrity	58
	3.11 Prolonged expression of GPI anchor-deficient GREG induces apoptosis	61
	3.12 Interaction between GREG and other GIC proteins	64
4	Discussions	67
	4.1 GREG, a GPI-anchored protein resident at the Golgi complex	68
	4.1.1 Golgi resident proteins	68
	4.1.2 Signals for Golgi targeting	70
	4.1.3 Requirement of the EQ repeats for Golgi localization of GREG	72
	4.1.4 Is the EQ repeat a universal Golgi targeting signal?	73
	4.1.5 Involvement of the SNARE-like domain in Golgi targeting	74
	4.1.6 Is oligomerization involved in Golgi localization of GREG	74
	4.2 Involvement of GREG in maintenance of the Golgi structure	77
	4.2.1 Is GREG a structural protein at the Golgi apparatus?	77
	4.2.2 Does GREG function via heterotrimeric G proteins?	78
	4.2.3 Alternative signaling pathways,	
	possibly mediating GREG function	81
	4.3 GREG as an essential component of Golgi-specific lipid rafts	82
	4.3.1 Biogenesis of GICs	82
	4.3.2 A model of GICs	83
	4.3.3 GICs as distinct microdomains with distinct function(s)	84
5	Materials and Methods	87
	5.1 Materials	87
	5.2 Methods	93
	5.2.1 Cell biological and Immunological methods	93
	5.2.1.1 Cell culture	93
	5 2 1 2 Transfection of cells	94

	Contents
5.2.1.3 Metabolic labeling of cells	95
5.2.1.4 Interference mediated by dsRNA	95
5.2.1.5 Immunofluorescence microscope	96
5.2.1.6 Peptide antibodies	97
5.2.1.6.1 Animal immunization	97
5.2.1.6.2 Affinity purification	97
5.2.1.7 Immunoprecipitation	98
5.2.2 Molecular biological methods	99
5.2.2.1 Extraction of total cellular RNA from	
monolayer cultured cells	99
5.2.2.2 Reverse transcription	100
5.2.2.3 Polymerase chain reaction	101
5.2.2.4 Subcloning	102
5.2.2.5 Phage display	103
5.2.2.5.1 Preparation of host cells	103
5.2.2.5.2 Titration of the cDNA library	103
5.2.2.5.3 Screening of the cDNA library	104
5.2.2.6 Northern blotting	106
5.2.2.7 In vitro transcription	106
5.2.2.8 In vitro translation	107
5.2.2.9 Overexpression of GREG in Pichia cells	107
5.2.2.9.1 Preparation of competent cells	107
5.2.2.9.2 Transformation of cells	108
5.2.2.9.3 Purification of over-expressed GREG proteins	108
5.2.3 Biochemical methods	109
5.2.3.1 Golgi preparation	109
5.2.3.2 Preparation of Golgi-derived lipid rafts	110
5.2.3.3 Quantification of proteins	111
5.2.3.4 Trypsin digestion	111
5.2.3.5 Glycosidase digestion	112
5.2.3.6 Detergent-phase separation using TX-114	112
5.2.3.7 Subcellular fractionation	113
5.2.3.8 Isolation of DRMs	114
5.2.3.9 Western blot	115

	Contents
5.2.3.9.1 Transfer proteins from an SDS-PAGE gel	
to a PVDF filter	115
5.2.3.9.2 Antibody incubation	115
5.2.3.10 Aerolysin overlay assay	116
5.2.4 Sequence analysis with the use of web-based programs	117
6 References	118
7 Acknowledgements	138